

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/106628/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Davies, William ORCID: <https://orcid.org/0000-0002-7714-2440> 2018. The steroid sulfate axis and its relationship to maternal behavior and mental health. *Journal of Molecular Endocrinology* 61 , T199-T210. 10.1530/JME-17-0219 file

Publishers page: <http://dx.doi.org/10.1530/JME-17-0219>
<<http://dx.doi.org/10.1530/JME-17-0219>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



The steroid sulfate axis and its relationship to maternal behavior and mental health

William Davies^{1,2,3*}

¹School of Psychology, Cardiff University, Cardiff, UK

²Medical Research Council Centre for Neuropsychiatric Genetics and Genomics and
Division of Psychological Medicine and Clinical Neurosciences, School of Medicine,
Cardiff University, UK

³Neuroscience and Mental Health Research Institute, Cardiff University, UK

*Correspondence address: School of Psychology, Tower Building, 70, Park Place,
Cardiff CF10 3AT, U.K. e-mail: daviesw4@cardiff.ac.uk Tel: +44-(0)29-2087-0152

Short title: Steroid sulfates and maternal mental health

Keywords: dehydroepiandrosterone sulfate, postpartum depression, postpartum
psychosis, steroid sulfatase, sulfotransferase

Main text word count: 5,257

Abstract

Steroid hormones can exist in functionally-dissociable sulfated and non-sulfated (free) forms and can exert profound effects on numerous aspects of mammalian physiology; the ratio of free to sulfated steroids is governed by the antagonistic actions of steroid sulfatase (STS) and sulfotransferase (SULT) enzymes. Here, I examine evidence from human and animal model studies which suggests that STS and its major substrate (dehydroepiandrosterone sulfate, DHEAS) and product (DHEA) can influence brain function, behavior and mental health, before summarising how the activity of this axis varies throughout mammalian pregnancy and the postpartum period. I then consider how the steroid sulfate axis might impact upon normal maternal behavior and how its dysfunction might contribute towards risk of postpartum psychiatric illness. Understanding the biological substrates underlying normal and abnormal maternal behavior will be important for maximising the wellbeing of new mothers and their offspring.

Word count: 140

An introduction to the steroid sulfate axis

Steroid hormones are synthesised within a number of endocrine body tissues (notably the adrenal gland, gonadal, breast, adipose and liver tissue in primates), and, as well as being utilised locally, may subsequently be transported elsewhere to elicit widespread physiological effects. The hydrophobicity of free steroids limits their ability to be transported within aqueous media; hence, transport of these compounds is facilitated by the addition of negatively-charged sulfate groups through esterification by the steroid sulfotransferase (SULT) enzymes SULT1E1, SULT2A1, or SULT2B1b, or, to a lesser extent, by SULT1A1 or SULT2B1a (Salman *et al.* 2011, Mueller *et al.* 2015). In addition to increasing the solubility of steroids, sulfation processes also increase their stability: circulating concentrations of sulfated steroids are typically substantially higher than circulating concentrations of their free steroid counterparts, and the former may act as 'reservoirs' for the peripheral formation of bioactive hormones (Mueller *et al.* 2015). Many common steroids can be sulfated including cholesterol, pregnenolone, estrone and dehydroepiandrosterone (DHEA) (Mueller *et al.* 2015).

Upon influx into cells in target tissues via organic anion transporter proteins, sulfated steroids are typically desulfated by hydrolysis to their unconjugated forms, which are generally considered to be more biologically-active and which can act as precursors for a variety of androgens and estrogens. Whilst multiple sulfotransferases can facilitate sulfation according to tissue-type, there is just one ubiquitous enzyme which cleaves sulfate groups from steroids: steroid sulfatase (STS, formerly known as arylsulfatase C). In the interests of clarity and brevity, in this review I focus upon the physiological roles of STS and dehydroepiandrosterone (DHEA), whose sulfated form (DHEAS) is the most abundant circulating steroid in humans (Neunzig & Bernhardt 2014).

77 **Steroid sulfatase: its regulation, expression and function**

78 Steroid sulfatase is encoded by the X-linked *STS* gene (Xp22.3). As the human *STS* gene
 79 escapes X-inactivation (Shapiro *et al.* 1979), and as its Y-linked paralogue is pseudogenic (Yen
 80 *et al.* 1988), expression of *STS* is 1-2fold higher in female than male tissues (including brain)
 81 during development and into adulthood, although whether this expression difference
 82 translates to significantly greater enzyme activity in female tissues is debatable (Cuevas-
 83 Covarrubias *et al.* 1993, Miranda-Duarte *et al.* 1999, Ugele & Regemann 2000, Nakamura *et*
 84 *al.* 2003, Steckelbroek *et al.* 2004, Kriz *et al.* 2008, HE O'Brien *et al.* manuscript in preparation);
 85 nevertheless, the possibility certainly exists that the physiological consequences of *STS*
 86 activity modulation could feasibly be more profound in women than in men. *STS* is expressed
 87 in a number of tissues many of which are involved in reproductive function, including: the
 88 placenta (highest expression), brain, ovary, mammary gland, testis, adipose tissue, thyroid
 89 gland and skin (Salido *et al.* 1990, Miki *et al.* 2002, Steckelbroek *et al.* 2004, Stergiakouli *et al.*
 90 2011, <https://www.ncbi.nlm.nih.gov/unigene> accessed 12th September 2017).

91 At the cellular level, the *STS* protein is largely located in the endoplasmic reticulum of
 92 the cell where it functions as a glycosylated homodimer; the catalytic activity of *STS* is
 93 dependent upon the presence of sulfatase-modifying factors (SUMFs) (Mueller *et al.* 2015).
 94 *STS* activity appears to be increased in response to stress/inflammation in various tissues with
 95 the gene being a target of NF- κ B; the resultant free steroid products, notably estrogens, may
 96 act as inflammation-suppressors (Dias & Selcer 2016, Jiang *et al.* 2016). *STS* has long been
 97 recognised as a therapeutic target in hormone-dependent cancers, and a number of effective
 98 and specific *STS* inhibitors have been developed which reduce the pool of androgens and
 99 estrogens in the vicinity of the cancer and which have potential clinical benefits (Purohit &
 100 Foster 2012).

The steroid sulfatase axis: its influence on brain function and behavior

Seminal work in rodents suggested that, in mammals, some steroids (and hence their sulfate conjugates) such as DHEA(S) could be synthesised *de novo* in the brain and may be regarded as neurosteroids (Corpechot *et al.* 1981). Subsequent work has shown that, whilst DHEA biosynthesis within the brain is possible, a second route by which DHEA appears in the brain in rodents (and feasibly humans too) is through the uptake, and subsequent rapid desulfation, of circulating DHEAS by organic ion transporters and steroid sulfatase respectively in the capillaries of the blood-brain barrier (Qaiser *et al.* 2017). Whilst the expression patterns of STS in the human blood-brain barrier have yet to be systematically assessed, in the developing human brain, STS is highly expressed throughout the thalamus with lower expression also seen in the olfactory epithelium, the cerebral cortex, the basal ganglia, the hypothalamus and pituitary gland, the choroid plexus and the cerebellar neuroepithelium (Stergiakouli *et al.* 2011); in adulthood, high levels of STS expression and associated enzyme activity persist in these regions (Perumal *et al.* 1973, Steckelbroeck *et al.* 2004, Kriz *et al.* 2008). Although STS activity in the brain is likely to have important and widespread developmental and ongoing effects (see later), sulfatase activity in this tissue is apparently dominated by sulfotransferase activity, and, in support of this idea, levels of sulfated steroids in the human brain (including those of DHEAS) may be relatively high (Maninger *et al.* 2009, Mueller *et al.* 2015).

At the molecular level, free and sulfated steroid hormones can modulate receptors influencing acute neuronal inhibition and excitation, as well as neurodevelopmental processes. For example, both DHEAS and pregnenolone sulfate act as antagonists at GABA_A receptors, as agonists at σ receptors, and as positive modulators at N-methyl-D-aspartic acid (NMDA) receptors (Reed *et al.* 2005, Qaiser *et al.* 2017). Importantly, the sulfated and unconjugated steroids may have differential effects and potencies e.g. DHEA has been

reported to act as a weaker GABA_A antagonist than DHEAS (Reed *et al.* 2005, Maninger *et al.* 2009). In addition to relatively weak agonistic effects at the androgen and estrogen receptors, DHEA and DHEAS may bind to, and activate, neurotrophin TrkA and p75^{NTR} receptors to attenuate neuronal apoptosis and hence influence neurodevelopment (Lazaridis *et al.* 2011).

Insights into the behavioral and brain processes that are mediated by STS and DHEA(S), potentially via the aforementioned molecular mechanisms, can be obtained by examining phenotypes in individuals in which: a) the *STS* gene is rendered non-functional (by natural or experimental means), b) the STS enzyme is acutely inhibited by a selective drug, or c) DHEA(S) has been administered. Less convincingly, it is possible to investigate the extent to which (peripheral) levels of DHEA(S) correlate with brain/ behavioral phenotypes of interest, and hence the extent to which changes in the former may cause the latter.

In humans, *STS* deficiency (arising from partial or complete deletion of the gene, or inactivating point mutations within it) results in the rare dermatological condition X-linked ichthyosis (XLI) (Fernandes *et al.* 2010). XLI chiefly affects males, and is associated with an elevated DHEAS/DHEA serum ratio, especially pre-pubertally (Idkowiack *et al.* 2016). Whilst there is currently little published literature on brain structure/function and biochemistry in individuals with XLI (Trent & Davies 2013), there is an emerging literature suggesting that boys with the condition may be at increased risk of developmental disorders such as Autism Spectrum Conditions (ASCs), Attention Deficit Hyperactivity Disorder (ADHD), and early-onset psychotic disorders (Kent *et al.* 2008, Chatterjee *et al.* 2016, Malik *et al.* 2017) whilst men with the condition may be at increased risk of both developmental and mood (unipolar depression and bipolar) disorders (Chatterjee *et al.* 2016). There is also some evidence that genetic variation within *STS* is associated with measures of attention in both clinical (ADHD) (Brookes *et al.* 2010, Stergiakouli *et al.* 2011, Wang *et al.* 2017) and healthy (Humby *et al.* 2017)

populations. These behavioral findings are consistent with the high expression of STS in brain regions involved in integrating and acting upon sensory information, and executive function.

Recapitulating the clinical findings in XLI, mice in which the *Sts* gene is deleted (or in which the STS enzyme is inhibited) show significantly reduced levels of serum DHEA and associated impairments in attention, altered response inhibition, hyperactivity, heightened emotional reactivity and aggression, and increased levels of behavioral perseveration (Davies *et al.* 2009, Trent *et al.* 2012b, Trent *et al.* 2013, Davies *et al.* 2014). Animal models allow the neurochemistry underlying behavioral abnormalities to be studied. *Sts* deletion in mice is associated with higher serotonin levels in the striatum and hippocampus (together with elevated hippocampal expression of the serotonin receptor 2c (*Htr2c*) gene), and reduced striatal noradrenaline turnover; the extent of serotonergic perturbation in *Sts*-deficient mice seems to correlate with the severity of some behavioral phenotypes (Trent *et al.* 2012a). Pharmacological studies in rats in which the STS enzyme was acutely inhibited have revealed changes in hippocampal acetylcholinergic release together with parallel changes in memory function (Rhodes *et al.* 1997, Babalola *et al.* 2012, Yue *et al.* 2016).

Experimental and correlational studies in animal models and human populations have linked altered DHEA(S) levels to a diverse and important range of behavioral phenotypes including: sexual function (Peixoto *et al.* 2017), aggression (Nicolas *et al.* 2001, Soma *et al.* 2015), locomotor activity (Strous *et al.* 2001, Trent *et al.* 2012b, Trent *et al.* 2013) and numerous aspects of mood and cognition (including attention) (Rhodes *et al.* 1996, Davies *et al.* 2009, Pluchino *et al.* 2015, Starka *et al.* 2015).

The data presented above establish that steroid sulfatase (and its dysfunction) can impact extensively upon normal brain function via multiple neural and neurochemical pathways; this action may be direct (i.e. within the brain itself), or, alternatively, may result

from extra-brain STS activity impacting upon the production and action of circulating levels of sulfated and free steroids including DHEA(S).

Changes in the steroid sulfate axis throughout mammalian pregnancy and the postpartum period

Throughout the childbearing process, women experience considerable hormonal fluctuations, including with regard to the steroid sulfate axis (Tal *et al.* 2000). However, longitudinal studies in which the levels, and sulfation status, of multiple hormones are measured across pregnancy and the postpartum period are scarce. Due to ethical and practical issues, most information in humans comes from analysis of peripheral tissues (blood, serum/ plasma, saliva and rarely cerebrospinal fluid) and therefore its relevance to the hormonal milieu experienced directly by the brain is questionable. Moreover, peripherally-detected maternal hormone levels may be influenced by multiple variable factors including: breastfeeding, stress exposure, use of recreational and therapeutic drugs, parity, maternal age and diet, and gender/number/size of the fetus(es), and understanding how unstable hormone levels relate to specific physiological phenotypes is therefore challenging. Whilst the use of neurobiologically-amenable mammalian animal models in which experimental variables can be controlled may circumvent these issues to some extent in *in vivo* systems, such models differ from humans in terms of both circulating hormone levels and reproductive traits such as number of offspring per pregnancy, or the extent and duration of postnatal maternal care; hence, extrapolating from models to man (or woman) should be done with caution.

In humans, from around nine weeks of pregnancy, a key role of the steroid sulfate axis is to generate precursors for the production of estrogens to be secreted into the maternal and fetal bloodstreams. Initially, sulfated C-19 steroids including DHEAS and 16 α -OH-DHEAS

produced by the maternal and fetal adrenal glands and fetal liver must undergo hydrolysis in the STS-rich syncytiotrophoblast of the placenta (Salido *et al.* 1990) before conversion by a series of enzymatic reactions to estrogens including estrone, estriol and estradiol; estrone, but not estradiol, is subsequently sulfated in the mother (Geyer *et al.* 2017).

As healthy pregnancies progress, there is a consistently-observed decrease in maternal DHEAS serum levels from non-pregnancy levels (apparently independent of fetal gender), perhaps as DHEAS is increasingly utilised for estrogen synthesis in the developing placenta; after parturition, maternal serum DHEAS levels rapidly rebound to pre-pregnancy levels (Tagawa *et al.* 2004, Soldin *et al.* 2005, Kuijper *et al.* 2013, Farrar *et al.* 2014). The data regarding systemic maternal DHEA levels throughout pregnancy and the postpartum period are less consistent. Some studies have demonstrated elevated serum/plasma DHEA levels during early-mid pregnancy, with a subsequent gradual decline up to one year postpartum (Nieschlag *et al.* 1974, Buckwalter *et al.* 1999, Tagawa *et al.* 2004); given DHEA's immunosuppressive effects and an increase in maternal cytokine markers after childbirth, this pattern of effects has been postulated to provide maximum protection for the incipient fetus from maternal immune surveillance (Tagawa *et al.* 2004). Other studies have suggested that peripheral DHEA levels are relatively unaffected by pregnancy and parturition (Buster *et al.* 1979, Soldin *et al.* 2005) or even that they increase across pregnancy and towards parturition in peripheral tissues (saliva or plasma) (Bird *et al.* 1980, Hampson *et al.* 2013). If DHEAS levels do fluctuate as outlined above during pregnancy/postpartum period, and DHEA levels remain in a comparatively steady state, then the DHEA/DHEAS ratio would be expected to be high during pregnancy and low during the postpartum period relative to values in non-pregnancy; in healthy populations where this ratio has been assessed longitudinally, this pattern of effects is indeed observed (Hill *et al.* 2002, Tagawa *et al.* 2004).

Presumably the above changes in DHEA/DHEAS ratio over the course of pregnancy and the postpartum period are related to the relative abundance and/or activity of the steroid sulfatase and sulfotransferase enzymes in cells contributing towards the hormonal milieu of the periphery. A main contributor to this ratio is the syncytiotrophoblast cells of the placenta, and expulsion of the STS-rich placenta after birth likely explains the rapid restoration of circulating maternal DHEAS levels. However, other tissues may also contribute: in healthy women, STS activity in leukocytes has been reported to be greater in third trimester pregnant women than in first trimester pregnant, or non-pregnant, women (Miyakawa *et al.* 1994), a finding consistent with the observed high DHEA/DHEAS ratio during late pregnancy. To the best of my knowledge, there has not yet been any systematic analysis of peripheral (leukocyte) STS activity throughout the postpartum period in humans.

Peripheral levels of sulfated and free steroids cannot provide reliable information on the activity of steroid sulfatase in the brain, and direct measurement of brain STS activity throughout pregnancy and the postpartum period in humans is currently unachievable. However, animal models, such as rodents, might provide some insights into human physiology (bearing in mind the caveats discussed above with respect to cross-species extrapolation). Mortaud and colleagues (1996) showed that, in whole female mouse brain, STS protein levels were more than two-fold higher in the lactating (postpartum) state relative to the pregnant (stage not specified) or non-pregnant state; whether this increase in protein level corresponded to an increase in enzyme activity in this state, or with brain DHEA(S) levels, was not assessed. Conversely, in rats, neither STS brain activity nor sulfotransferase liver activity appear to be affected by pregnancy or parturition although only cortical (as opposed to whole) brain tissue was analysed (Maayan *et al.* 2004a). Interestingly, data on STS activity in rat leukocytes partially resemble those seen in humans, in that activity is significantly higher

in late-pregnancy animals (18 days post conception) than in non-pregnant animals, and activity becoming even more pronounced 24hrs after giving birth (Maayan *et al.* 2004a). In rat serum and brain cortex, the DHEA/DHEAS ratio is significantly, and equivalently, elevated in late pregnancy and early postpartum animals compared to non-pregnant control females (Maayan *et al.* 2004a).

In summary, the sparse human and rodent data presented above are reconcilable with the proposal that in non-cortical regions of the mammalian brain, and in certain cells of the immune system, STS levels/activity increase over the course of pregnancy before peaking during late pregnancy and into the early postpartum period.

A possible role for the steroid sulfate axis in normal maternal behavior

Androgen-related metabolic pathways, including the STS/DHEA(S) axis, are known to modulate physiological processes associated with parturition (Makieva *et al.* 2014). Given the previously-described role of the steroid sulfatase axis in brain and behavioral function, its increased activity in the perinatal period may be related to, and potentially be causal for, the emergence of maternal behaviors designed to nourish and protect their offspring. These behaviors, many of which are highly-conserved across mammalian species, include: nest-building, huddling, nursing and social interaction with the offspring mediated by olfactory, visual, auditory and somatosensory cues, altered (generally decreased) levels of anxiety with increased exploratory behavior, and aggression directed towards threatening predators/society members but not offspring (Bridges 2015, Lonstein *et al.* 2015). In rodents, and probably also in humans, the quality and intensity of expressed maternal behaviors is related to maternal cognitive (executive) function, particularly in the domains of offspring-related learning and memory processes, attention to relevant care cues, behavioral flexibility and impulse regulation; interestingly, in rats, reduced maternal behavior is associated with

impaired performance on attentional set-shifting and startle/prepulse inhibition tasks, whilst increased maternal care (pup-licking) is related to reduced motor impulsivity (Lonstein *et al.* 2015). In non-primate species, and in primates to a lesser extent, these behaviors are driven by hormonal mechanisms acting via a multitude of brain regions and neurochemical systems, including the prefrontal cortex, the amygdala, the cholinergic basal forebrain activating system and the GABAergic and serotonergic systems (Bridges 2015, Lonstein *et al.* 2015). The fact that manipulation of the STS axis in males affects many of the cognitive/behavioral phenotypes and neurobiological systems listed above (notably attention, social interaction, emotional reactivity, aggression, memory, behavioral flexibility and motor impulsivity (Mortaud *et al.* 1996, Rhodes *et al.* 1997, Kent *et al.* 2008, Davies *et al.* 2009, Trent *et al.*, 2012a,2012b, Trent *et al.* 2013, Davies *et al.* 2014, Chatterjee *et al.* 2016)) supports the argument that the STS axis influences neural processes pertinent to maternal care efficacy in females. To explicitly test the idea that the STS axis influences maternal perinatal behavior, studies will need to be undertaken in non-pregnant (control), pregnant and postpartum female animal models and human subjects in which STS activity is compromised, or in which DHEA(S) levels are systematically varied and assayed.

To date, the only available *Sts*-deficient genetic rodent models have been chromosomally-mutant mice that are necessarily male (Trent *et al.* 2012b), but new gene editing technology should hopefully allow the generation of *Sts*-deficient female rodents (Baud & Flint 2017). The expectation that such genetically-modified female rodents may exhibit STS-dependent abnormal maternal behaviors has been raised by a recent pharmacological study in our laboratory. Briefly, we showed that acute inhibition of STS in new mouse mothers resulted in anxiety-related phenotypes (a reduced startle response, and increased rearing and exploratory drive on the elevated plus maze), but no gross

abnormalities in nest maintenance or in mother-pup interactions (Humby *et al.* 2016). At this stage, we cannot discount the fact that there were subtle, undetectable, irregularities in dam-pup interactions, especially in light of the fact that inhibitor-treated mothers exhibit substantial dysregulation of olfactory-related gene expression in the brain (W Davies, unpublished results). Furthermore, as we did not examine the behavioral effects of acute STS inhibition in female mice with other physiological statuses (virgin, non-virgin but non-pregnant, and pregnant), we cannot definitively say that the behavioral effects mediated by the STS axis are specific to the postpartum period; this extended analysis is ongoing. We are currently undertaking a parallel systematic study of behavior, including perinatal behavior, in STS-deficient women with a view to determining which, if any, psychological processes are affected by their genetic mutation.

Additional evidence that DHEA(S) participate in normal perinatal maternal behaviors in humans may be obtained by showing behavioral effects elicited by administration to new mothers, or through identifying significant correlations between systemic DHEA(S) levels and behavioral/cognitive measures in healthy (or general population) postpartum mothers. Although DHEAS administration has been performed in postpartum women (e.g. Aisaka *et al.* 1984) there is no reliable published data available regarding parallel behavioral changes. In healthy women selected from the general population, there is some evidence for an association between higher serum DHEA levels, enhanced mood and aspects of better cognitive performance during late pregnancy (~20 days prior to delivery) and the postpartum period (~26 days after delivery) (Buckwalter *et al.* 1999), although no similar relationship seems to exist for DHEAS levels (Farrar *et al.* 2014).

Postpartum psychiatric disorders

A significant proportion of women experience mental health issues manifesting in late pregnancy and/or in the postpartum period. These can range from relatively common and comparatively mild conditions which do not require medical intervention (so-called ‘baby blues’) to rarer, more severe, persistent and disabling disorders which require urgent medical care; the latter category of disorders includes postpartum depression, obsessive-compulsive disorder and anxiety disorders (Sharma & Sommerdyk 2015, Stewart & Vigod 2016, Pawluski *et al.* 2017). If the steroid sulfate axis is indeed a major player in maternal brain and behavioral function in late pregnancy and the postpartum period as has been argued above, then, logically, its dysfunction might reasonably be considered a risk factor for vulnerability to maternal mental health conditions in this period. In the following section, I discuss the nature and aetiology of one extremely severe and poorly-understood psychiatric disorder associated with childbirth, postpartum (or puerperal) psychosis, and consider evidence implicating steroid sulfate axis abnormalities in its pathogenesis.

The nature and etiology of postpartum psychosis

Postpartum psychosis (PP) is estimated to affect 1-2 in every 1000 new mothers (VanderKruik *et al.* 2017). Symptoms associated with the condition include hallucinations, delusions (often related to the newborn child), cognitive disorganisation, anxiety, and mood abnormalities, and these tend to present within the first two weeks (and frequently within the first few days) of childbirth; PP symptoms can impact massively upon normal mother-child bonding and family life, and affected mothers are at elevated risk of committing suicide or infanticide (Bergink *et al.* 2016). Whilst there is thought to be a considerable biological component to disorder vulnerability, the exact nature and contribution of underlying biological risk factors is currently obscure (Jones *et al.* 2014, Bergink *et al.* 2016). Understanding these may help to

develop better predictive biomarkers for the condition, as well as more effective and safer treatment options (Davies 2017). Epidemiological data has suggested considerable overlap with bipolar disorder, autoimmune thyroid conditions, and pre-eclampsia (Jones *et al.* 2014, Bergink *et al.* 2016); other studies have implicated an abnormal (over-active) immune system (Bergink *et al.* 2013, Kumar *et al.* 2017), autoimmune anti-NMDA receptor encephalitis (Bergink *et al.* 2015), and serotonergic system dysfunction (Kumar *et al.* 2007, Davies 2017) in risk. A genetic linkage study in bipolar postpartum psychosis implicated significant and suggestive loci at 16p13 and 8q24 respectively (Jones *et al.* 2007), but, to date, findings from small-scale (and therefore underpowered) candidate gene and genome-wide association studies have been unconvincing or non-significant, and none have yet implicated X-linked genetic risk variants.

Steroid sulfate axis dysfunction and postpartum psychosis risk

There are several lines of basic and clinical evidence suggesting that abnormalities with the steroid sulfate axis (and most likely steroid sulfatase deficiency (Davies 2012)) may influence PP risk: a) the axis appears to exert disproportionately large effects in the late pregnancy/early postpartum period, so any disruption to it may impact relatively specifically on this timepoint, b) estrogens are generally thought to be protective against psychosis (Reicher-Rossler 2017) and STS deficiency, in women, as in men (Lykkesfeldt *et al.* 1985), is expected to result in lower levels of circulating estrogens as a consequence of reduced levels of DHEA precursor, c) genetic deletions encompassing *STS* have been associated with psychotic disorders (paranoid and early-onset schizophrenia) in case studies (Milunsky *et al.* 1999, Malik *et al.* 2017), d) *STS* brain expression is high in regions previously implicated in psychotic disorders (Fusar-Poli *et al.* 2011, Dietsche *et al.* 2017), e) the neurochemical abnormalities associated with psychosis and remediable by antipsychotic treatment overlap

365 considerably with those influenced by STS and DHEA(S) i.e. of the serotonergic system
 366 (notably the 5-HT_{2c} receptor)(Meltzer *et al.* 2012, Selvaraj *et al.* 2014), the hippocampal
 367 cholinergic system (Olincy & Freedman 2012, Carruthers *et al.* 2015), the GABA_A system
 368 (Egerton *et al.* 2017), and of NMDA receptor signalling (notably in the thalamus)(Vukadinovic
 369 2014, Harrison 2015), f) STS-deficient humans and mice exhibit a range of PP-relevant
 370 phenotypes including inattention and emotional instability, whilst genetic variants within *STS*
 371 in man are associated with cognitive disorganisation (see above), g) STS is highly-expressed in
 372 the hypothalamus, pituitary gland and the thyroid gland (Stergiakouli *et al.* 2011) and its
 373 absence in these tissues could potentially explain high rates of hypothalamus-pituitary-
 374 thyroid axis dysfunction and autoimmune thyroid dysfunction in PP, h) abnormal placental
 375 and whole blood STS expression is associated with pre-eclampsia (Gratton *et al.* 2016), i)
 376 levels of salivary DHEAS during late pregnancy and in the early postpartum period (10 days
 377 after birth) positively correlate with measures of anxiety, phobia, paranoia and psychoticism
 378 in previously-healthy women, with highest DHEAS levels (consistent with impaired or absent
 379 STS activity) being associated with significant psychiatric distress (Marrs *et al.* 2009, Marrs *et*
 380 *al.* 2010), j) lithium, an established effective treatment for mania in bipolar disorder and PP,
 381 enhances the serum DHEA/DHEAS ratio in rats consistent with a stimulatory effect on STS,
 382 whilst reducing both brain and serum levels of DHEAS (Maayan *et al.* 2004b), k) pathologically-
 383 reduced levels of immunosuppressive DHEA in the postpartum period as a consequence of
 384 STS deficiency may feasibly contribute towards the immune hyper-activation seen in PP and
 385 l) *STS* and the *HTR2C* (5-HT_{2c}) gene lie under candidate quantitative trait loci linkage peaks in
 386 a porcine model of PP (Quilter *et al.* 2007). Finally, the prevalence of STS deficiency i.e.
 387 heterozygosity or homozygosity for null mutations in women is estimated to be ~1 in 950
 388 individuals (based upon the general population frequency of STS deficiency in males (Langlois

et al. 2009, Craig *et al.* 2010) and *de novo* versus inherited mutation rates (Cuevas-Covarrubias *et al.* 1999)); this rate is consistent with it being a risk factor for PP, although clearly, as for most mood and psychotic disorders, multiple interacting genetic and environmental risk factors are likely to influence overall PP vulnerability.

Future analyses, in STS-deficient women or in *Sts*-deficient female rodents, are likely to provide evidence for or against the hypothesis that this molecular perturbation increases PP risk, although such studies will likely be limited by available sample size, by the infrequency of the condition, and by cross-species extrapolation issues. Further studies in healthy women in late pregnancy and the postpartum period, which will be less constrained by sample size, might examine if, and how, peripheral steroid sulfatase activity (in addition to DHEA(S) levels) correlate with dimensional behavioral measures related to PP (e.g. psychoticism). Alternative complementary work in clinical PP populations might investigate: a) variability within the *STS* gene (where rates of causal polymorphisms/mutations might be expected to differ from control women), b) STS activity in accessible tissues such as leukocytes (lower activity anticipated in women affected by PP than in controls) or c) peripheral baseline and stress-evoked levels of DHEA(S)(higher DHEAS/DHEA ratios expected in affected women versus control subjects).

Our mouse studies have provided preliminary evidence somewhat supportive of the notion of STS deficiency as a risk factor for PP. Briefly, we found that the behavioral phenotypes elicited by STS inhibition in new mothers could be partially reversed by administration of the clinically-efficacious antipsychotic drug ziprasidone, thus indicating their potential relevance to PP (Humby *et al.* 2016). Additionally, and intriguingly, STS-inhibited mice demonstrated abnormal gene expression within a small region of chromosome 15 (equivalent to the 8q24 candidate genomic region implicated by linkage in PP), which could

also be normalised by ziprasidone administration, providing further support for the model's face and predictive validity and thus its utility for understanding the mechanistic basis of PP risk (Humby *et al.* 2016).

Steroid sulfate axis dysfunction in other postpartum psychiatric disorders

Postpartum psychosis is an umbrella term covering a wide variety of behavioral and psychiatric symptoms, in addition to psychosis, that present shortly after childbirth. Many of these symptoms (which can include depressive and manic episodes, anxiety and obsessive-compulsive tendencies) may also be seen, to varying extents, in cases of other differentially-defined postpartum psychiatric conditions such as postpartum depression, and could be underpinned by common biological processes. Much of the logic implicating steroid sulfate axis dysfunction in general, and STS deficiency in particular, in PP pathophysiology may be equally applied to these alternative disorders, especially considering that their occurrence may, in part, be due to abnormalities in physiological and neurochemical systems affected by this axis i.e. steroid hormone levels, the hypothalamic-pituitary-adrenal axis, the thyroid system, markers of inflammation and the GABAergic, noradrenergic and serotonergic systems (Speisman *et al.* 2011, Skalkidou *et al.* 2012, Pawluski *et al.* 2017). The observed effects of STS deficiency on mood modulation (Chatterjee *et al.* 2016), behavioral (in)flexibility (Trent *et al.* 2013) and anxiety-related processes (Trent *et al.* 2012b, Chatterjee *et al.* 2016) implicate it as a candidate risk factor in postpartum depression, OCD, and anxiety respectively. The steroid sulfate axis could also feasibly mediate the effects of environmental factors (e.g. stressors such as childhood maltreatment) on mothers' vulnerability to a range of postpartum psychiatric illnesses, and potentially even on their infants health (Sexton *et al.* 2015, Schury *et al.* 2017).

Evidence examining rates of these various postpartum psychiatric conditions and associated behavioral and physiological markers remains to be collected in STS-deficient individuals and animal models; conversely, rates of STS deficiency remain to be determined in patients ascertained on the basis of having been diagnosed with such postpartum conditions. However, there is some limited existing data, consistent with the STS deficiency risk hypothesis, suggesting that circulating maternal DHEA plasma levels may be abnormally low prior to, and during the onset of, postpartum depression (Gelman *et al.* 2015), and that DHEA supplementation can benefit mood in depressed individuals (although consistent therapeutic benefits of this intervention in the postpartum period have yet to demonstrated)(Soares & Phillips 2006, Peixoto *et al.* 2014).

Conclusions and future work

Above I have discussed the mounting evidence (admittedly mainly obtained in male test subjects to date) that steroid sulfatase, and the reactions it catalyses, have important roles in a wide variety of important brain and behavioral functions. As the activity of the STS axis fluctuates across numerous tissues (including brain) during pregnancy, and into the postpartum period, it is conceivable that this axis bears upon normal maternal behavioral phenotypes and its influence in this respect remains to be tested using genetic and pharmacological approaches in clinical and model populations. In particular, gene expression changes elicited by manipulations of the steroid sulfate axis may be compared against those seen in the healthy postpartum maternal brain (Gammie *et al.* 2016). It also follows that, potentially, STS axis dysfunction may be associated with postpartum psychiatric conditions, and there is some circumstantial evidence, notably in postpartum psychosis, that this may be the case. Genetic and endocrine analyses in women previously affected by, or at high of

developing, postpartum psychiatric conditions should be able to experimentally test this hypothesis.

Of course, whilst I have focussed upon the possible impact of STS and DHEA(S) on maternal behavior here, it is naïve to think that these molecules act in isolation to affect this phenotype. As such, future studies should aim to supplement the currently available, very limited, data relating to the brain and peripheral expression and activity of enzymes and compounds involved in DHEA(S) biosynthesis and metabolism (notably the SULTs and SUMFs) across pregnancy and the postpartum period in healthy women, women affected by postpartum psychiatric illness, and relevant mammalian animal models, with a view to understanding how these components interact to generate healthy or abnormal behaviours. Work in model systems in particular may highlight molecules and pathways which mediate the behavioral effects of the steroid sulfate axis, and which could theoretically comprise novel therapeutic targets. For example, our nascent mouse work has implicated the CCN family of proteins as potentially-druggable targets in cases of PP (Davies, 2017). Pharmacological or genetic targeting of such mediators might result in lower levels of side-effects compared to the systemic administration of compounds such as DHEA or estrogens which have androgenic/estrogenic potential and which elicit widespread effects on multiple physiological systems (Gentile 2005, Rutkowski *et al.* 2014).

Declaration of interest

There is no conflict of interest that could be perceived as prejudicing the impartiality of this review.

Funding

This work was undertaken within the Medical Research Council UK Centre for Neuropsychiatric Genetics and Genomics (MR/L010305/1) but did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

References

- Aisaka K, Mori H, Ogawa T & Kigawa T 1984 Effects of dehydroepiandrosterone-sulphate (DHEA-S) administration on puerperal lactation and maternal prolactin and estradiol levels. *Nihon Sanka Fujinka Gakkai Zasshi* **36** 1935-1942.
- Babalola PA, Fitz NF, Gibbs RB, Flaherty PT, Li PK & Johnson DA 2012 The effect of the steroid sulfatase inhibitor (p-O-sulfamoyl)-tetradecanoyl tyramine (DU-14) on learning and memory in rats with selective lesion of septal-hippocampal cholinergic tract. *Neurobiol Learn Mem* **98** 303-310.
- Baud A & Flint J 2017 Identifying genes for neurobehavioral traits in rodents: progress and pitfalls. *Dis Model Mech* **10** 373-383.
- Bergink V, Armangue T, Titulaer MJ, Markx S, Dalmau J & Kushner SA 2015 Autoimmune Encephalitis in Postpartum Psychosis. *Am J Psychiatry* **172** 901-908.
- Bergink V, Burgerhout KM, Weigelt K, Pop VJ, de Wit H, Drexhage RC, Kushner SA & Drexhage HA 2013 Immune system dysregulation in first-onset postpartum psychosis. *Biol Psychiatry* **73** 1000-1007.
- Bergink V, Rasgon N & Wisner KL 2016 Postpartum Psychosis: Madness, Mania, and Melancholia in Motherhood. *Am J Psychiatry* **173** 1179-1188.
- Bird CE, Cook SJ, Dressel DE & Clark AF 1980 Plasma C19-delta 5-steroid levels during normal human pregnancy. *Clin Biochem* **13** 126-128.
- Bridges RS 2015 Neuroendocrine regulation of maternal behavior. *Front Neuroendocrinol* **36** 178-196.
- Brookes KJ, Hawi Z, Park J, Scott S, Gill M & Kent L 2010 Polymorphisms of the steroid sulfatase (STS) gene are associated with attention deficit hyperactivity disorder and influence brain tissue mRNA expression. *Am J Med Genet B Neuropsychiatr Genet* **153B** 1417-1424.

537 Buckwalter JG, Stanczyk FZ, McCleary CA, Bluestein BW, Buckwalter DK, Rankin KP, Chang L &
 538 Goodwin TM 1999 Pregnancy, the postpartum, and steroid hormones: effects on cognition
 539 and mood. *Psychoneuroendocrinology* **24** 69-84.
 540 Buster JE, Chang RJ, Preston DL, Elashoff RM, Cousins LM, Abraham GE, Hobel CJ & Marshall
 541 JR 1979 Interrelationships of circulating maternal steroid concentrations in third trimester
 542 pregnancies. II. C18 and C19 steroids: estradiol, estriol, dehydroepiandrosterone,
 543 dehydroepiandrosterone sulfate, delta 5-androstenediol, delta 4-androstenedione,
 544 testosterone, and dihydrotestosterone. *J Clin Endocrinol Metab* **48** 139-142.
 545 Carruthers SP, Gurvich CT & Rossell SL 2015 The muscarinic system, cognition and
 546 schizophrenia. *Neurosci Biobehav Rev* **55** 393-402.
 547 Chatterjee S, Humby T & Davies W 2016 Behavioral and Psychiatric Phenotypes in Men and
 548 Boys with X-Linked Ichthyosis: Evidence from a Worldwide Online Survey. *PLoS One* **11**
 549 e0164417.
 550 Corpechot C, Robel P, Axelson M, Sjoval J & Baulieu EE 1981 Characterization and
 551 measurement of dehydroepiandrosterone sulfate in rat brain. *Proc Natl Acad Sci U S A* **78**
 552 4704-4707.
 553 Craig WY, Roberson M, Palomaki GE, Shackleton CH, Marcos J & Haddow JE 2010 Prevalence
 554 of steroid sulfatase deficiency in California according to race and ethnicity. *Prenat Diagn* **30**
 555 893-898.
 556 Cuevas-Covarrubias SA, Juarez-Oropeza MA, Miranda-Zamora R & Diaz-Zagoya JC 1993
 557 Comparative analysis of human steroid sulfatase activity in prepubertal and postpubertal
 558 males and females. *Biochem Mol Biol Int* **30** 691-695.

559 Cuevas-Covarrubias SA, Valdes-Flores M, Orozco Orozco E, Diaz-Zagoya JC & Kofman-Alfaro
 560 SH 1999 Most "sporadic" cases of X-linked ichthyosis are not de novo mutations. *Acta Derm*
 561 *Venereol* **79** 143-144.
 562 Davies W 2012 Does steroid sulfatase deficiency influence postpartum psychosis risk? *Trends*
 563 *Mol Med* **18** 256-262.
 564 Davies W 2017 Understanding the pathophysiology of postpartum psychosis: Challenges and
 565 new approaches. *World J Psychiatry* **7** 77-88.
 566 Davies W, Humby T, Kong W, Otter T, Burgoyne PS & Wilkinson LS 2009 Converging
 567 pharmacological and genetic evidence indicates a role for steroid sulfatase in attention. *Biol*
 568 *Psychiatry* **66** 360-367.
 569 Davies W, Humby T, Trent S, Eddy JB, Ojarikre OA & Wilkinson LS 2014 Genetic and
 570 pharmacological modulation of the steroid sulfatase axis improves response control;
 571 comparison with drugs used in ADHD. *Neuropsychopharmacology* **39** 2622-2632.
 572 Dias NJ & Selcer KW 2016 Steroid sulfatase in the human MG-63 preosteoblastic cell line:
 573 Antagonistic regulation by glucocorticoids and NFkappaB. *Mol Cell Endocrinol* **420** 85-96.
 574 Dietsche B, Kircher T & Falkenberg I 2017 Structural brain changes in schizophrenia at
 575 different stages of the illness: A selective review of longitudinal magnetic resonance imaging
 576 studies. *Aust N Z J Psychiatry* **51** 500-508.
 577 Egerton A, Modinos G, Ferrera D & McGuire P 2017 Neuroimaging studies of GABA in
 578 schizophrenia: a systematic review with meta-analysis. *Transl Psychiatry* **7** e1147.
 579 Farrar D, Tuffnell D, Neill J, Scally A & Marshall K 2014 Assessment of cognitive function across
 580 pregnancy using CANTAB: a longitudinal study. *Brain Cogn* **84** 76-84.
 581 Fernandes NF, Janniger CK & Schwartz RA 2010 X-linked ichthyosis: an oculocutaneous
 582 genodermatosis. *J Am Acad Dermatol* **62** 480-485.

583 Fusar-Poli P, Borgwardt S, Crescini A, Deste G, Kempton MJ, Lawrie S, Mc Guire P & Sacchetti
 584 E 2011 Neuroanatomy of vulnerability to psychosis: a voxel-based meta-analysis. *Neurosci*
 585 *Biobehav Rev* **35** 1175-1185.

586 Gammie SC, Driessen TM, Zhao C, Saul MC & Eisinger BE 2016 Genetic and neuroendocrine
 587 regulation of the postpartum brain. *Front Neuroendocrinol* **42** 1-17.

588 Gelman PL, Flores-Ramos M, Lopez-Martinez M, Fuentes CC & Grajeda JP 2015 Hypothalamic-
 589 pituitary-adrenal axis function during perinatal depression. *Neurosci Bull* **31** 338-350.

590 Gentile S 2005 The role of estrogen therapy in postpartum psychiatric disorders: an update.
 591 *CNS Spectr* **10** 944-952.

592 Geyer J, Bakhaus K, Bernhardt R, Blaschka C, Dezhkam Y, Fietz D, Grosser G, Hartmann K,
 593 Hartmann MF, Neunzig J, et al. 2017 The role of sulfated steroid hormones in reproductive
 594 processes. *J Steroid Biochem Mol Biol* **172** 207-221.

595 Gratton AM, Ye L, Brownfoot FC, Hannan NJ, Whitehead C, Cannon P, Deo M, Fuller PJ, Tong
 596 S & Kaitu'u-Lino TJ 2016 Steroid sulfatase is increased in the placentas and whole blood of
 597 women with early-onset preeclampsia. *Placenta* **48** 72-79.

598 Hampson E, Phillips SD, Soares CN & Steiner M 2013 Steroid concentrations in antepartum
 599 and postpartum saliva: normative values in women and correlations with serum. *Biol Sex*
 600 *Differ* **4** 7.

601 Harrison PJ 2015 Recent genetic findings in schizophrenia and their therapeutic relevance. *J*
 602 *Psychopharmacol* **29** 85-96.

603 Hill M, Parizek A, Klak J, Hampl R, Sulcova J, Havlikova H, Lapcik O, Bicikova M, Fait T, Kancheva
 604 R, et al. 2002 Neuroactive steroids, their precursors and polar conjugates during parturition
 605 and postpartum in maternal and umbilical blood: 3.3beta-hydroxy-5-ene steroids. *J Steroid*
 606 *Biochem Mol Biol* **82** 241-250.

607 Humby T, Cross ES, Messer L, Guerrero S & Davies W 2016 A pharmacological mouse model
 608 suggests a novel risk pathway for postpartum psychosis. *Psychoneuroendocrinology* **74** 363-
 609 370.

610 Humby T, Fisher A, Allen C, Reynolds M, Hartman A, Giegling I, Rujescu D & Davies W 2017 A
 611 genetic variant within STS previously associated with inattention in boys with attention deficit
 612 hyperactivity disorder is associated with enhanced cognition in healthy adult males. *Brain*
 613 *Behav* **7** e00646.

614 Idkowiak J, Taylor AE, Subtil S, O'Neil DM, Vijzelaar R, Dias RP, Amin R, Barrett TG, Shackleton
 615 CH, Kirk JM, et al. 2016 Steroid Sulfatase Deficiency and Androgen Activation Before and After
 616 Puberty. *J Clin Endocrinol Metab* **101** 2545-2553.

617 Jiang M, Klein M, Zanger UM, Mohammad MK, Cave MC, Gaikwad NW, Dias NJ, Selcer KW,
 618 Guo Y, He J, et al. 2016 Inflammatory regulation of steroid sulfatase: A novel mechanism to
 619 control estrogen homeostasis and inflammation in chronic liver disease. *J Hepatol* **64** 44-52.

620 Jones I, Chandra PS, Dazzan P & Howard LM 2014 Bipolar disorder, affective psychosis, and
 621 schizophrenia in pregnancy and the post-partum period. *Lancet* **384** 1789-1799.

622 Jones I, Hamshere M, Nangle JM, Bennett P, Green E, Heron J, Segurado R, Lambert D,
 623 Holmans P, Corvin A, et al. 2007 Bipolar affective puerperal psychosis: genome-wide
 624 significant evidence for linkage to chromosome 16. *Am J Psychiatry* **164** 1099-1104.

625 Kent L, Emerton J, Bhadravathi V, Weisblatt E, Pasco G, Willatt LR, McMahon R & Yates JR
 626 2008 X-linked ichthyosis (steroid sulfatase deficiency) is associated with increased risk of
 627 attention deficit hyperactivity disorder, autism and social communication deficits. *J Med*
 628 *Genet* **45** 519-524.

629 Kriz L, Bicikova M, Mohapl M, Hill M, Cerny I & Hampl R 2008 Steroid sulfatase and sulfuryl
 630 transferase activities in human brain tumors. *J Steroid Biochem Mol Biol* **109** 31-39.

631 Kuijper EA, Ket JC, Caanen MR & Lambalk CB 2013 Reproductive hormone concentrations in
 632 pregnancy and neonates: a systematic review. *Reprod Biomed Online* **27** 33-63.
 633 Kumar HB, Purushottam M, Kubendran S, Gayathri P, Mukherjee O, Murthy AR, Ghosh S,
 634 Chandra P, Reddy YC, Benegal V, et al. 2007 Serotonergic candidate genes and puerperal
 635 psychosis: an association study. *Psychiatr Genet* **17** 253-260.
 636 Kumar MM, Venkataswamy MM, Sathyanarayanan G, Thippeswamy H, Chandra PS & Mani
 637 RS 2017 Immune system aberrations in postpartum psychosis: An immunophenotyping study
 638 from a tertiary care neuropsychiatric hospital in India. *J Neuroimmunol* **310** 8-13.
 639 Langlois S, Armstrong L, Gall K, Hulait G, Livingston J, Nelson T, Power P, Pugash D, Siciliano
 640 D, Steinraths M, et al. 2009 Steroid sulfatase deficiency and contiguous gene deletion
 641 syndrome amongst pregnant patients with low serum unconjugated estriols. *Prenat Diagn* **29**
 642 966-974.
 643 Lazaridis I, Charalampopoulos I, Alexaki VI, Avlonitis N, Pediaditakis I, Efstathopoulos P,
 644 Calogeropoulou T, Castanas E & Gravanis A 2011 Neurosteroid dehydroepiandrosterone
 645 interacts with nerve growth factor (NGF) receptors, preventing neuronal apoptosis. *PLoS Biol*
 646 **9** e1001051.
 647 Lonstein JS, Levy F & Fleming AS 2015 Common and divergent psychobiological mechanisms
 648 underlying maternal behaviors in non-human and human mammals. *Horm Behav* **73** 156-185.
 649 Lykkesfeldt G, Bennett P, Lykkesfeldt AE, Micic S, Moller S & Svenstrup B 1985 Abnormal
 650 androgen and estrogen metabolism in men with steroid sulphatase deficiency and recessive
 651 X-linked ichthyosis. *Clin Endocrinol (Oxf)* **23** 385-393.
 652 Maayan R, Abou-Kaud M, Strous RD, Kaplan B, Fisch B, Shinnar N & Weizman A 2004a The
 653 influence of parturition on the level and synthesis of sulfated and free neurosteroids in rats.
 654 *Neuropsychobiology* **49** 17-23.

- 655 Maayan R, Shaltiel G, Poyurovsky M, Ramadan E, Morad O, Nechmad A, Weizman A & Agam
 656 G 2004b Chronic lithium treatment affects rat brain and serum dehydroepiandrosterone
 657 (DHEA) and DHEA-sulphate (DHEA-S) levels. *Int J Neuropsychopharmacol* **7** 71-75.
- 658 Makieva S, Saunders PT & Norman JE 2014 Androgens in pregnancy: roles in parturition. *Hum*
 659 *Reprod Update* **20** 542-559.
- 660 Malik A, Amer AB, Salama M, Haddad B, Alrifai MT, Al Balwi M, Davies W & Eyaid W 2017 X-
 661 linked ichthyosis associated with psychosis and behavioral abnormalities: a case report.
 662 *Journal of Medical Case Reports*.
- 663 Maninger N, Wolkowitz OM, Reus VI, Epel ES & Mellon SH 2009 Neurobiological and
 664 neuropsychiatric effects of dehydroepiandrosterone (DHEA) and DHEA sulfate (DHEAS). *Front*
 665 *Neuroendocrinol* **30** 65-91.
- 666 Marrs CR, Ferraro DP & Cross CL 2010 Hormones and mood across the first postpartum year.
 667 *Eur J Obstet Gynecol Reprod Biol* **149** 121-122.
- 668 Marrs CR, Ferraro DP, Cross CL & Rogers SL 2009 A potential role for adrenal androgens in
 669 postpartum psychiatric distress. *Eur J Obstet Gynecol Reprod Biol* **143** 127-128.
- 670 Meltzer HY, Massey BW & Horiguchi M 2012 Serotonin receptors as targets for drugs useful
 671 to treat psychosis and cognitive impairment in schizophrenia. *Curr Pharm Biotechnol* **13** 1572-
 672 1586.
- 673 Miki Y, Nakata T, Suzuki T, Darnel AD, Moriya T, Kaneko C, Hidaka K, Shiotsu Y, Kusaka H &
 674 Sasano H 2002 Systemic distribution of steroid sulfatase and estrogen sulfotransferase in
 675 human adult and fetal tissues. *J Clin Endocrinol Metab* **87** 5760-5768.
- 676 Milunsky J, Huang XL, Wyandt HE & Milunsky A 1999 Schizophrenia susceptibility gene locus
 677 at Xp22.3. *Clin Genet* **55** 455-460.

678 Miranda-Duarte A, Valdes-Flores M, Miranda-Zamora R, Diaz-Zagoya JC, Kofman-Alfaro SH &
679 Cuevas-Covarrubias SA 1999 Steroid sulfatase activity in leukocytes: a comparative study in
680 45,X; 46,Xi(Xq) and carriers of steroid sulfatase deficiency. *Biochem Mol Biol Int* **47** 137-142.
681 Miyakawa I, Kawano Y, Taniyama K & Mori N 1994 Steroid sulfatase activity in human
682 leukocytes. *Gynecol Obstet Invest* **38** 191-193.
683 Mortaud S, Donsez-Darcel E, Roubertoux PL & Degrelle H 1996 Murine steroid sulfatase gene
684 expression in the brain during postnatal development and adulthood. *Neurosci Lett* **215** 145-
685 148.
686 Mueller JW, Gilligan LC, Idkowiak J, Arlt W & Foster PA 2015 The Regulation of Steroid Action
687 by Sulfation and Desulfation. *Endocr Rev* **36** 526-563.
688 Nakamura Y, Miki Y, Suzuki T, Nakata T, Darnel AD, Moriya T, Tazawa C, Saito H, Ishibashi T,
689 Takahashi S, et al. 2003 Steroid sulfatase and estrogen sulfotransferase in the atherosclerotic
690 human aorta. *Am J Pathol* **163** 1329-1339.
691 Neunzig J & Bernhardt R 2014 Dehydroepiandrosterone sulfate (DHEAS) stimulates the first
692 step in the biosynthesis of steroid hormones. *PLoS One* **9** e89727.
693 Nicolas LB & Fry JP 2007 The steroid sulfatase inhibitor COUMATE attenuates rather than
694 enhances access of dehydroepiandrosterone sulfate to the brain in the mouse. *Brain Res* **1174**
695 92-96.
696 Nieschlag E, Walk T & Schindler AE 1974 Dehydroepiandrosterone (DHA) and DHA-sulphate
697 during pregnancy in maternal blood. *Horm Metab Res* **6** 170-171.
698 Olincy A & Freedman R 2012 Nicotinic mechanisms in the treatment of psychotic disorders: a
699 focus on the alpha7 nicotinic receptor. *Handb Exp Pharmacol* 211-232.
700 Pawluski JL, Lonstein JS & Fleming AS 2017 The Neurobiology of Postpartum Anxiety and
701 Depression. *Trends Neurosci* **40** 106-120.

- 702 Peixoto C, Carrilho CG, Barros JA, Ribeiro TT, Silva LM, Nardi AE, Cardoso A & Veras AB 2017
 703 The effects of dehydroepiandrosterone on sexual function: a systematic review. *Climacteric*
 704 **20** 129-137.
- 705 Peixoto C, Devicari Cheda JN, Nardi AE, Veras AB & Cardoso A 2014 The effects of
 706 dehydroepiandrosterone (DHEA) in the treatment of depression and depressive symptoms in
 707 other psychiatric and medical illnesses: a systematic review. *Curr Drug Targets* **15** 901-914.
- 708 Perumal AS & Robins E 1973 Regional and subcellular distribution of aryl-and steroid
 709 sulfatases in brain. *Brain Res* **59** 349-358.
- 710 Pluchino N, Drakopoulos P, Bianchi-Demicheli F, Wenger JM, Petignat P & Genazzani AR 2015
 711 Neurobiology of DHEA and effects on sexuality, mood and cognition. *J Steroid Biochem Mol*
 712 *Biol* **145** 273-280.
- 713 Purohit A & Foster PA 2012 Steroid sulfatase inhibitors for estrogen- and androgen-
 714 dependent cancers. *J Endocrinol* **212** 99-110.
- 715 Qaiser MZ, Dolman DEM, Begley DJ, Abbott NJ, Cazacu-Davidescu M, Corol DI & Fry JP 2017
 716 Uptake and metabolism of sulphated steroids by the blood-brain barrier in the adult male rat.
 717 *J Neurochem* **142** 672-685.
- 718 Quilter CR, Blott SC, Wilson AE, Bagga MR, Sargent CA, Oliver GL, Southwood OI, Gilbert CL,
 719 Mileham A & Affara NA 2007 Porcine maternal infanticide as a model for puerperal psychosis.
 720 *Am J Med Genet B Neuropsychiatr Genet* **144B** 862-868.
- 721 Reed MJ, Purohit A, Woo LW, Newman SP & Potter BV 2005 Steroid sulfatase: molecular
 722 biology, regulation, and inhibition. *Endocr Rev* **26** 171-202.
- 723 Rhodes ME, Li PK, Burke AM & Johnson DA 1997 Enhanced plasma DHEAS, brain acetylcholine
 724 and memory mediated by steroid sulfatase inhibition. *Brain Res* **773** 28-32.

- 725 Rhodes ME, Li PK, Flood JF & Johnson DA 1996 Enhancement of hippocampal acetylcholine
726 release by the neurosteroid dehydroepiandrosterone sulfate: an in vivo microdialysis study.
727 *Brain Res* **733** 284-286.
- 728 Riecher-Rossler A 2017 Estrogens, prolactin, hypothalamic-pituitary-gonadal axis, and
729 schizophrenic psychoses. *Lancet Psychiatry* **4** 63-72.
- 730 Rutkowski K, Sowa P, Rutkowska-Talipska J, Kuryliszyn-Moskal A & Rutkowski R 2014
731 Dehydroepiandrosterone (DHEA): hypes and hopes. *Drugs* **74** 1195-1207.
- 732 Salido EC, Yen PH, Barajas L & Shapiro LJ 1990 Steroid sulfatase expression in human placenta:
733 immunocytochemistry and in situ hybridization study. *J Clin Endocrinol Metab* **70** 1564-1567.
- 734 Salman ED, Faye-Petersen O & Falany CN 2011 Hydroxysteroid sulfotransferase 2B1b
735 expression and localization in normal human brain. *Horm Mol Biol Clin Investig* **8** 445-454.
- 736 Schury K, Koenig AM, Isele D, Hulbert AL, Krause S, Umlauf M, Kolassa S, Ziegenhain U,
737 Karabatsiakakis A, Reister F, et al. 2017 Alterations of hair cortisol and dehydroepiandrosterone
738 in mother-infant-dyads with maternal childhood maltreatment. *BMC Psychiatry* **17** 213.
- 739 Selvaraj S, Arnone D, Cappai A & Howes O 2014 Alterations in the serotonin system in
740 schizophrenia: a systematic review and meta-analysis of postmortem and molecular imaging
741 studies. *Neurosci Biobehav Rev* **45** 233-245.
- 742 Sexton MB, Hamilton L, McGinnis EW, Rosenblum KL & Muzik M 2015 The roles of resilience
743 and childhood trauma history: main and moderating effects on postpartum maternal mental
744 health and functioning. *J Affect Disord* **174** 562-568.
- 745 Shapiro LJ, Mohandas T, Weiss R & Romeo G 1979 Non-inactivation of an x-chromosome locus
746 in man. *Science* **204** 1224-1226.
- 747 Sharma V & Sommerdyk C 2015 Obsessive-compulsive disorder in the postpartum period:
748 diagnosis, differential diagnosis and management. *Womens Health (Lond)* **11** 543-552.

- 749 Skalkidou A, Hellgren C, Comasco E, Sylven S & Sundstrom Poromaa I 2012 Biological aspects
750 of postpartum depression. *Womens Health (Lond)* **8** 659-672.
- 751 Soares CN & Phillips SD 2006 Hormones in the treatment of depression. *Womens Health*
752 *(Lond)* **2** 435-445.
- 753 Soldin OP, Guo T, Weiderpass E, Tractenberg RE, Hilakivi-Clarke L & Soldin SJ 2005 Steroid
754 hormone levels in pregnancy and 1 year postpartum using isotope dilution tandem mass
755 spectrometry. *Fertil Steril* **84** 701-710.
- 756 Soma KK, Rendon NM, Boonstra R, Albers HE & Demas GE 2015 DHEA effects on brain and
757 behavior: insights from comparative studies of aggression. *J Steroid Biochem Mol Biol* **145**
758 261-272.
- 759 Speisman BB, Storch EA & Abramowitz JS 2011 Postpartum obsessive-compulsive disorder. *J*
760 *Obstet Gynecol Neonatal Nurs* **40** 680-690.
- 761 Starka L, Duskova M & Hill M 2015 Dehydroepiandrosterone: a neuroactive steroid. *J Steroid*
762 *Biochem Mol Biol* **145** 254-260.
- 763 Stergiakouli E, Langley K, Williams H, Walters J, Williams NM, Suren S, Giegling I, Wilkinson LS,
764 Owen MJ, O'Donovan MC, et al. 2011 Steroid sulfatase is a potential modifier of cognition in
765 attention deficit hyperactivity disorder. *Genes Brain Behav* **10** 334-344.
- 766 Stewart DE & Vigod S 2016 Postpartum Depression. *N Engl J Med* **375** 2177-2186.
- 767 Strous RD, Spivak B, Yoran-Hegesh R, Maayan R, Averbuch E, Kotler M, Mester R & Weizman
768 A 2001 Analysis of neurosteroid levels in attention deficit hyperactivity disorder. *Int J*
769 *Neuropsychopharmacol* **4** 259-264.
- 770 Tagawa N, Hidaka Y, Takano T, Shimaoka Y, Kobayashi Y & Amino N 2004 Serum
771 concentrations of dehydroepiandrosterone and dehydroepiandrosterone sulfate and their

772 relation to cytokine production during and after normal pregnancy. *Clin Chim Acta* **340** 187-
 773 193.

774 Tal R, Taylor HS, Burney RO, Mooney SB & Giudice LC 2000 Endocrinology of Pregnancy. In
 775 *Endotext*. Eds LJ De Groot, G Chrousos, K Dungan, KR Feingold, A Grossman, JM Hershman, C
 776 Koch, M Korbonits, R McLachlan, M New, et al. South Dartmouth (MA).

777 Trent S, Cassano T, Bedse G, Ojarikre OA, Humby T & Davies W 2012a Altered serotonergic
 778 function may partially account for behavioral endophenotypes in steroid sulfatase-deficient
 779 mice. *Neuropsychopharmacology* **37** 1267-1274.

780 Trent S & Davies W 2013 Cognitive, behavioral and psychiatric phenotypes associated with
 781 steroid sulfatase deficiency. *World Journal of Translational Medicine* **2** 1-12.

782 Trent S, Dean R, Veit B, Cassano T, Bedse G, Ojarikre OA, Humby T & Davies W 2013 Biological
 783 mechanisms associated with increased perseveration and hyperactivity in a genetic mouse
 784 model of neurodevelopmental disorder. *Psychoneuroendocrinology* **38** 1370-1380.

785 Trent S, Dennehy A, Richardson H, Ojarikre OA, Burgoyne PS, Humby T & Davies W 2012b
 786 Steroid sulfatase-deficient mice exhibit endophenotypes relevant to attention deficit
 787 hyperactivity disorder. *Psychoneuroendocrinology* **37** 221-229.

788 Ugele B & Regemann K 2000 Differential increase of steroid sulfatase activity in XX and XY
 789 trophoblast cells from human term placenta with syncytia formation in vitro. *Cytogenet Cell*
 790 *Genet* **90** 40-46.

791 VanderKruik R, Barreix M, Chou D, Allen T, Say L, Cohen LS & Maternal Morbidity Working G
 792 2017 The global prevalence of postpartum psychosis: a systematic review. *BMC Psychiatry* **17**
 793 272.

794 Vukadinovic Z 2014 NMDA receptor hypofunction and the thalamus in schizophrenia. *Physiol*
 795 *Behav* **131** 156-159.

- 796 Wang LJ, Chan WC, Chou MC, Chou WJ, Lee MJ, Lee SY, Lin PY, Yang YH & Yen CF 2017
797 Polymorphisms of STS gene and SULT2A1 gene and neurosteroid levels in Han Chinese boys
798 with attention-deficit/hyperactivity disorder: an exploratory investigation. *Sci Rep* **7** 45595.
- 799 Yen PH, Marsh B, Allen E, Tsai SP, Ellison J, Connolly L, Neiswanger K & Shapiro LJ 1988 The
800 human X-linked steroid sulfatase gene and a Y-encoded pseudogene: evidence for an
801 inversion of the Y chromosome during primate evolution. *Cell* **55** 1123-1135.
- 802 Yue XH, Tong JQ, Wang ZJ, Zhang J, Liu X, Liu XJ, Cai HY & Qi JS 2016 Steroid sulfatase inhibitor
803 DU-14 protects spatial memory and synaptic plasticity from disruption by amyloid beta
804 protein in male rats. *Horm Behav* **83** 83-92.